

Avoid micronutrient depletion

Growers run into major problems when they don't pay attention to their micros

By Carl Whitcomb

Recently, a nurseryman told me he had inadvertently left out Micromax from a batch of mix last growing season and the plants grew off fine.

Based on this incident, he was seeking my advice before stopping the addition of micronutrients entirely to save money.

My advice was to continue to add 1.5 pounds of Micromax micronutrients per cubic yard of mix. The pennies saved could cost him dearly down the road. It's important to understand what happened to the plants that received no Micromax and why adding micronutrients is important to continued good plant growth.

Micronutrients (iron, manganese, copper, boron, zinc and molybdenum — and perhaps chlorine, nickel and cobalt in miniscule amounts) are appropriately named in that the amount needed by the plant relative to nitrogen, phosphorus, potassium, calcium, magnesium and sulfur is indeed quite small, or micro, in proportion.

But, the role micronutrients play is critical to continued good plant growth.

Auto example

Being an old car buff, I sometimes describe the nutrient elements relative to aspects of an automobile. For example, nitrogen is like gasoline in the tank. More is required than anything else and, by adding more, growth is faster.

Phosphorus is like oil in the crankcase — an ingredient critical to the production and transfer of energy, and so on. I describe molybdenum like the small wire that connects the coil with the distributor. It's just a small piece of wire, typically no more than 8-12 inches long, and miniscule relative to everything else. But without it you walk.

When I began my research into the complex of growing plants in containers at the University of Florida in 1968, micronutrient deficiencies were rampant. Some very dramatic plant responses to micronutrient treatments occurred during that time.



Cutting-grown plants may have residual micronutrients from parent plants stored in leaf and stem tissues.

For several studies I obtained liners from commercial nurseries and placed them in experiments where varying levels and proportions of micronutrients had been added. The plants would go from having off-colored and sometimes chlorotic leaves and dull leaf surfaces to lots of bud breaks and vibrant new growth.

I recall thinking, "Wow, I have found the answer." But when cuttings were taken from the best plants in the study, rooted and placed again into the same or similar nutritional

treatment as before, plant response was good, but was not as dramatic as with the parents.

In some cases chlorosis or other micronutrient deficiency symptoms gradually developed. What happened?

Plant reactions

Plant growth is most affected by the most deficient element or complex of deficient elements. The more severe the deficiency, the greater plant response when the deficient element is added.

But with cuttings taken from

those plants, deficiencies were less severe, so plant response was less dramatic. Also, it is important to realize that even with the plant response to the micronutrient elements added, the levels and proportions were far from being correct, as would be revealed in later studies.

In looking back at some of those early studies, it is likely that plants went from a deficiency of one micronutrient element to an excess, thus further complicating the sorting out of cause and effect.

Another factor that came into play, and added to the intrigue, was the fact that on most occasions cuttings taken from plants growing in the field were less responsive to micronutrient treatments compared to cuttings taken from plants growing in containers.

I began to wonder about the residual of micronutrient elements in plant tissue. In other words, the levels and proportions of micronutrients in a good field soil could provide certain concentrations in stems and leaves of cuttings. If so, how long would it take for those micronutrient levels to decline to the point of causing deficiency symptoms?

Copper study

To study this, I chose copper as the element to vary, primarily because visual symptoms of copper deficiency are rather distinct. I took cuttings from Chinese holly plants growing in a good field soil with suitable micronutrient levels and rooted those cuttings in a propagation mix with no micronutrients added.

The liners were planted into a container growth medium with no copper added, but the other micronutrients were added (the best levels I knew at the time). After one growing season the plants looked normal in every way so cuttings were taken from those plants and rooted in a propagation mix with no micronutrients added.

The following growing season the liners were again planted onto a container growth medium with no copper added but the other micronutrients present. At the end of the second growing season the plants still looked normal. Again, cuttings were taken from those plants and rooted and the procedure repeated.

During the third growing season symptoms typical of copper deficiency began to appear early and by mid season symptoms were dramatic and crop growth and quality was poor.

**SUCCESS IS
IN THE ROOTS.**

Long-lasting Perennials that bloom the first year.

You have to dig deep to find Perennials as well-grounded as Speedling's first-year bloomers. From a rich history in growing, our vegetative and seed-propagated plants burst to life with robust, well-established root systems you can finish and "turn" with confidence. With a multitude of new and classic varieties, your customers will be Perennially happier. And you thought our success was an Annual thing.

SPEEDLING
ORNAMENTALS

800.426.4400 www.speedling.com

▲ Request 63 ▲

It Works!

Phyton-27® delivers exceptional bacterial and fungal disease control on a wide range of ornamental crops. Preventively and therapeutically, from propagation to post-harvest, it just works!

Savvy growers rely on *Phyton-27*!

Phyton
27
BACTERICIDE & FUNGICIDE

NEW
California
Registration

Phyton-27®
New
Dimension
BACTERICIDE & FUNGICIDE

Introducing **Phyton-27®**

New Dimension

- CAUTION signal word
- Easier to mix & clean up
- Same reliable disease control



Available in
1 & 2.5
gallon
bottles



For technical information, call 1-800-356-8733 or visit our website at www.phytoncorp.com
Phyton Corporation - Manufacturer of Phyton-27® Read and follow all label directions.

▲ Request 77 ▲



Moisty says, "TRY MOISTURIN"

A safe non-toxic spray that forms a clear impermeable film on plants.

BLOCKS OUT INSECTS & DISEASE
LOCKS IN MOISTURE DURING STRESS

GSI Horticultural



Bend, Oregon 541-383-0222
www.gsihorticultural.com

▲ Request 78 ▲

Copper deficiency symptoms are exceptionally dark green leaves, reduced leaf size, death of terminal growth and a proliferation of buds on lower stems.

To continue this study, cuttings were taken from the plants now showing copper deficiency and rooted. Cuttings were also taken from the original parent plants growing in field soil and rooted.

The following growing season, both sources of liners were planted into containers where the growth medium had the best complex of all micronutrient elements known at the time.

Liners from the copper deficient parents produced dramatic new growth free of deficiency symptoms. By the end of the growing season, plants from the severely deficient parents were similar in size, branching and overall quality as plants from parents grown in field soil and that showed no copper deficiency.

Cuttings were taken from plants with both treatments, rooted and grown in containers with all micronutrients added the following year. There were no detectable differences in plant response or plant quality the second year.

What to learn

The take-home message from this series of experiments is that copper deficiency does not occur abruptly. It was not until the third production cycle — with no copper added to the growth medium of the parent or the offspring's — when tissue levels in the plant were diluted to the point where growth was dramatically affected and copper deficiency symptoms were evident.

Copper levels in recently matured leaves were monitored during the studies. Interestingly, tissue analysis varied more between individual plants with the same treatment compared to between plants with or without copper added to the growth medium and provided little useful information.

Based on this and other work, I am not a fan of tissue analysis for diagnosing micronutrient deficiencies, with a few exceptions.

Birch problems

Micronutrient depletion also appears to play a role in river birch. River birch trees grown in field soils rarely show the stunting and abnormal leaf development referred to as



Mouse-ear disease on river birch, shown on the right, has been shown to be a nickel deficiency. It doesn't occur with field-grown material.

mouse-ear disease.

This is a problem unique to container-grown plants and date back only to about 1980. Container growth media components such as pine bark, peat, sand or other materials typically contain virtually no nickel. Micronutrient fertilizers do not include nickel and would have only miniscule amounts as impurities.

With refinements in slow-release fertilizers, single or triple superphosphate (which contained significant nickel) are rarely added to container growth media anymore. Applications of nickel sulfate have been shown to cause the symptoms to disappear. It appears that mouse-ear disease of river birch results from the near complete exclusion of nickel as an impurity in fertilizer components and dilution of nickel in plant tissue. On the other hand, plants grown in the field have sufficient nickel available for normal growth.

Plants grown from seeds have typically been more sensitive to micronutrient deficiencies vs. plants grown from cuttings. This is probably due to the larger amount of micronutrients residual in cutting tissue compared to endosperm in seeds.

Since my research that created Micromax micronutrient fertilizer in 1980, deficiencies are rare where nurseries consistently add 1.5 pounds of Micromax per cubic yard of growth medium and use good, quality irrigation water. Adding 1.5 pounds of Micromax per cubic yard is inexpensive insurance against tissue micronutrient depletion.

Carl Whitcomb is founder of Lacebark Inc., 2104 N. Cottonwood Road, Stillwater, OK 74075-2098; (405) 377-3539; fax (405) 377-0131; www.lacebarkinc.com.

**For More Information
Contact Our
Grower Technical Sales Office
(888) 447-3403**

bwi
Companies, Inc.

GTS

▲ Request 88 ▲

Offering Superior American Native Trees

Native Trees

Woody Warehouse NURSERY INC.

Specializing in native trees and shrubs grown with the latest root-pruning technology to yield massive fibrous roots and eliminate root circling.

Customized contract growing services for seedlings to caliper sized trees.

Available this Fall
Sturdy liners in 5" cinder bags
Ask about our larger sizes

Now in full **Root Force™** with **RootMakers®**

Woody Warehouse Nursery Inc. ph 317-994-5487
P.O. Box 259 fax 317-994-5494
Lizton, IN 46149-0259 toll free 866-766-8367
www.woodywarehouse.com

CONTRACT GROWING • LINERS • LANDSCAPE •
MITIGATION • REFORESTATION • WILDLIFE PLANTINGS

▲ Request 89 ▲